

CHANGING FREQUENCY SEPARATION OF KILOHERTZ QUASI-PERIODIC OSCILLATIONS IN THE SONIC-POINT BEAT-FREQUENCY MODEL

FREDERICK K. LAMB^{1,2,3} AND M. COLEMAN MILLER⁴

Submitted July 29, 2000

ABSTRACT

Previous work on the sonic-point beat-frequency (SPBF) model of the kilohertz quasi-periodic oscillations (QPOs) observed in the X-ray flux from neutron stars in low-mass binary systems has shown that it naturally explains many properties of these QPOs. These include the existence of just two principal QPOs in a given source, the commensurability of the frequency separation $\Delta\nu$ of the two kilohertz QPOs and the spin frequency ν_s inferred from burst oscillations, and the high frequencies, coherence, and amplitudes of these QPOs. Here we show that the SPBF model predicts that $\Delta\nu$ is less than but close to ν_s , consistent with the observed differences between $\Delta\nu$ and ν_s . It also explains naturally the decrease in $\Delta\nu$ with increasing QPO frequency seen in some sources and the plateau in the QPO frequency–X-ray flux observed in 4U 1820–30. The model fits well the QPO frequency behavior observed in Sco X-1, 4U 1608–52, 4U 1728–34, and 4U 1820–30 ($\chi^2/\text{dof} = 0.4\text{--}2.1$), giving masses ranging from 1.59 to 2.0 $M_\odot$ and spin rates ranging from 279 to 364 Hz. In the SPBF model, the kilohertz QPOs are effects of strong-field gravity. Thus, if the model is validated, the kilohertz QPOs can be used not only to determine the properties of neutron stars but also to explore quantitatively general relativistic effects in the strong-field regime.

Subject headings: relativity: general — stars: neutron — stars: oscillations — stars: rotation — stars: X-ray

1. INTRODUCTION

Kilohertz quasi-periodic brightness oscillations (QPOs) have been discovered in the accretion-powered emission of more than twenty neutron star low-mass X-ray binaries using the *Rossi X-ray Timing Explorer (RXTE)* (see van der Klis 2000 for a review). These kilohertz QPOs have frequencies ranging from ~ 500 Hz up to ~ 1200 Hz, amplitudes $\sim 1\%$ – 15% rms in the 2–60 keV band of the *RXTE* Proportional Counter Array, and quality factors $Q \equiv \nu/\text{FWHM}$ as large as $\sim 100\text{--}200$. They often appear as two simultaneous peaks in power spectra of the X-ray brightness, at frequencies ν_1 and ν_2 ($> \nu_1$) that move together and can shift upward and downward in frequency by up to a factor of two within a few hundred seconds, apparently as a consequence of changes in the accretion rate. The frequencies of the kilohertz QPOs lie within the range expected for orbital motion near neutron stars and are almost certainly strong-field general relativistic phenomenon (see Lamb, Miller, & Psaltis 1998; Miller 2000; van der Klis 2000).

In six of the kilohertz QPO sources, strong (amplitudes up to 50% rms) brightness oscillations with frequencies of hundreds of Hertz have been observed during thermonuclear X-ray bursts. There is very strong evidence that these oscillations are produced by rotation of brighter regions of the stellar surface with the spin of the star and that the most prominent oscillation frequency is the stellar spin frequency ν_s or its first overtone (see, e.g., Strohmayer & Markwardt 1999; Miller 1999). The coherent oscillations observed in burst tails, after hot matter has spread around the star, and the antipodal brighter regions observed in 4U 1636–53 are strong evidence that these stars have dynamically important surface magnetic fields (Miller 1999), as expected on evolutionary grounds (Miller, Lamb, &

Psaltis 1998, hereafter MLP98; Lorimer 2000). In the four sources where burst oscillations and two simultaneous kilohertz QPOs have both been detected with high confidence, the difference $\Delta\nu \equiv \nu_2 - \nu_1$ between the frequencies of the kilohertz QPOs agrees with the fundamental burst oscillation frequency to within 0.7% to 15% (see van der Klis 2000).

The properties of the kilohertz QPOs strongly indicate that the upper kilohertz QPO is an orbital frequency and that the lower is generated by the beat of this frequency with the star's spin (Lamb et al. 1998; MLP98; Miller, Lamb, & Cook 1998): (1) The frequencies of the kilohertz QPOs are in the range expected for orbital motion near neutron stars, consistent with one being an orbital frequency. (2) The relatively small variation of $\Delta\nu$ in a given source and the approximate commensurability of $\Delta\nu$ with the stellar spin frequency inferred from burst oscillations indicates that the spin of the star is generating the frequency difference. (3) The observation of at most two strong kilohertz QPOs in any given source (van der Klis 2000; Méndez & van der Klis 2000), rather than three, indicates that only a single sideband is being generated, rather than the two sidebands generated by other mechanisms. (4) In order to beat with the stellar spin, the other motion must be rotation about the center of the neutron star, indicating that the other motion is orbital motion. (5) Accretion of angular momentum ensures that the star is spinning in the prograde direction, implying that the frequency ν_2 of the upper kilohertz QPO is close to the orbital frequency ν_{orb} and that the frequency ν_1 of the lower kilohertz QPO is close to the beat frequency $\nu_B \equiv \nu_{\text{orb}} - \nu_s$.

This and other evidence motivated the development of the sonic-point beat-frequency (SPBF) model (MLP98). In this model ν_2 is close to the orbital frequency ν_{orb} at the sonic ra-

¹Center for Theoretical Astrophysics, University of Illinois at Urbana-Champaign, 1110 W. Green Street, Urbana, IL 61801-3080.

²Department of Physics, University of Illinois at Urbana-Champaign.

³Department of Astronomy, University of Illinois at Urbana-Champaign.

⁴Department of Astronomy, University of Maryland at College Park, College Park, MD 20742-2421.

dius, where the flow in the disk changes from nearly circular to rapidly inspiraling, and ν_1 is comparable to ν_B . The transition to hypersonic radial inflow is an effect of strong-field gravity. In the SPBF model, the low-frequency QPOs seen in the Z and atoll sources are produced by the magnetospheric beat-frequency mechanism (Alpar & Shaham 1985; Lamb et al. 1985). The model explains naturally many features of the kilohertz QPOs in addition to their frequencies and frequency separation and the existence of only two principal QPOs in a given source (MLP98). In the SPBF model, the kilohertz QPOs are a strong-field general relativistic effect and are therefore sensitive probes of the properties of the spacetime near the neutron star, including whether there is an innermost stable circular orbit (ISCO) and the gravitomagnetic torque created by the spin of the neutron star.

Careful analyses have shown that in some kilohertz QPO sources $\Delta\nu$ decreases systematically by 30–100 Hz, depending on the source, as ν_2 increases by a much larger amount (Sco X-1: van der Klis et al. 1997; 4U 1608–52: Méndez et al. 1998; 4U 1735–44: Ford et al. 1998; 4U 1728–34: Méndez & van der Klis 1999; see also Psaltis et al. 1998) and that the separation frequency in 4U 1636–536 is slightly but significantly smaller than the spin frequency inferred from its burst oscillations (Méndez, van der Klis, & van Paradijs 1998).

Here we show that in the SPBF model, the inward radial velocity of the accretion flow increases ν_1 and decreases ν_2 , making $\Delta\nu$ smaller than ν_s . An increase in ν_1 of $\lesssim 5\%$ is sufficient to account for the largest observed differences between $\Delta\nu$ and ν_B and decreases of $\Delta\nu$ with increasing ν_2 . The inward radial velocity of the accretion flow was included in the gas dynamical and radiation transport calculations reported by MLP98, but the effect of this motion on the frequencies of the kilohertz QPOs was neglected. In § 2 we discuss the generation of the kilohertz QPOs in the SPBF model, derive general expressions for their frequencies, taking into account the inward motion of the accretion flow, summarize the results of numerical simulations, and introduce the model of the gas dynamics in the SPBF model that we use here to calculate the frequencies of the kilohertz QPOs. In § 3 we show that despite its simplicity, this model fits accurately the frequency behavior observed in Sco X-1, 4U 1608–52, 4U 1728–34, and 4U 1820–30 and gives reasonable masses and spin rates for these neutron stars.

2. KILOHERTZ QPOS IN THE SONIC-POINT MODEL

Generation of kilohertz QPOs.—In the SPBF model, the kilohertz QPOs are produced by clumping of gas in the accretion flow near the star, probably caused by interaction of the accreting gas with the weak stellar magnetic field (see Ghosh & Lamb 1979; Aly & Kuipers 1990). Only clumps near the sonic radius R_{sp} , where the inward radial velocity of gas leaving the clumps becomes supersonic, contribute to the oscillations (see MLP98). At low accretion rates, radiation drag removes angular momentum from the gas in the clumps, causing it to spiral inward to the stellar surface. At high enough accretion rates, the optical depth from the stellar surface is large and radiation drag is unimportant. If, however, the star is smaller than the ISCO, gas will fall inward supersonically from clumps near the ISCO, where general relativity causes gas to spiral inward even if it does not lose any angular momentum. In either case, the supersonic inflow is a strong-field general relativistic effect.

Where gas falling inward from clumps near R_{sp} impacts the stellar surface, it produces bright footprints that move around the star with a frequency nearly equal to the orbital frequency

ν_{orb} of the clumps at R_{sp} , causing the X-ray flux seen by distant observers to oscillate as the footprint moves into and out of view. (As discussed by MLP98, ν_{orb} is not exactly equal to the Keplerian frequency ν_K , because the radiation force has a radial component and the clumps are drifting slowly inward.) *This oscillation is the upper kilohertz QPO; its frequency ν_2 is determined by the rate at which the azimuthal phase of the footprint advances.*

The weak magnetic field of the star funnels extra gas toward the magnetic poles, producing a broad beam of extra radiation that rotates with the star. Clumps orbiting near R_{sp} move through this beam with frequency $\nu_B \equiv \nu_{\text{orb}} - \nu_s$. When a clump is illuminated by the beam, gas in it loses angular momentum to the radiation at a faster rate, so more gas falls inward, causing the luminosity of the footprint to oscillate with a frequency close to ν_B . *This oscillation is the lower kilohertz QPO; its frequency ν_1 is determined by the frequency at which the mass flux onto a footprint oscillates.*

The flux in both oscillations is generated primarily at the stellar surface, where most of the gravitational energy is released, and hence both oscillations can have relatively high amplitudes.

Kilohertz QPO frequencies and frequency separation.—The kilohertz QPO frequencies predicted by the SPBF model can be calculated as follows (we use Boyer-Lindquist coordinates for easy reference to infinity). The light travel time from the stellar surface to the sonic radius is very small and hence the time between two successive maxima of the mass inflow rate from a given clump is very close to the beat period $\Delta t_B \equiv 1/(\nu_{\text{orb}} - \nu_s)$. Let t_1 and $t_2 = t_1 + \Delta t_B$ be the times of successive maxima, let r_1 and r_2 and $2\pi\phi_1$ and $2\pi\phi_2$ be the radial and azimuthal coordinates of the clump at t_1 and t_2 , let Δt_1 and Δt_2 be the times required for the gas stripped from the clump at t_1 and t_2 to reach the surface of the star, and let $2\pi\Delta\phi_1$ and $2\pi\Delta\phi_2$ be the changes in the azimuthal phase of the gas released at t_1 and t_2 as it falls from the clump to the stellar surface. Then the frequencies of the lower and upper kilohertz QPOs are

$$\nu_1 = \frac{1}{t_2 - t_1 + (\Delta t_2 - \Delta t_1)} \quad (1)$$

and

$$\nu_2 = \frac{\phi_2 - \phi_1 + (\Delta\phi_2 - \Delta\phi_1)}{t_2 - t_1 + (\Delta t_2 - \Delta t_1)}. \quad (2)$$

Suppose first that the inward motion of the clumps near R_{sp} can be neglected. This is the approximation used in MLP98 to compute the frequencies of the kilohertz QPOs. Then the time required for gas to spiral inward from a given clump to the stellar surface and the change in the azimuthal phase of the gas during its inspiral do not depend on when the gas separated from the clump, i.e., $\Delta t_2 = \Delta t_1$ and $\Delta\phi_2 = \Delta\phi_1$. Because $\Delta t_2 = \Delta t_1$, the time between two successive maxima of the mass flow rate at the stellar surface is the same as at the clump, where it is Δt_B , and hence the frequency ν_1 of the luminosity oscillation is $1/\Delta t_B = \nu_{\text{orb}} - \nu_s \equiv \nu_B$ (see eq. [1]). Because the rate at which the azimuthal phase of the footprint advances is the same as the rate at which the azimuthal phase of the clump advances, the rotation frequency ν_2 of the footprint is $(\phi_2 - \phi_1)/(t_2 - t_1) = \nu_{\text{orb}}$ (see eq. [2]) and the difference between ν_2 and ν_1 is therefore ν_s .

Suppose now that the inward motion of the clumps near R_{sp} is too large to be neglected. The radial distance of a given clump decreases with time and therefore the time required for gas to

spiral inward from the clump to the stellar surface steadily decreases. Hence $\Delta t_2 < \Delta t_1$. Equation (1) shows that to first order in v_{cl} , the frequency ν_1 of the lower kilohertz QPO is $\nu_B(1 + v_{cl}\partial_r\Delta t) > \nu_B$; ν_1 is greater than ν_B because the inward motion of the clump reduces the spatial separation between successive maxima of the mass flow rate, analogous to the Doppler shift of a sound wave. Equation (2) shows that to first order in v_{cl} , the frequency ν_2 of the upper kilohertz QPO is $\nu_{orb}(1 + v_{cl}\partial_r\Delta t) - v_{cl}\partial_r\Delta\phi - \nu_B^{-1}v_{cl}\partial_r\nu_{orb}$. Numerical simulations of the gas flow and radiation transport in full general relativity similar to those presented in MLP98 show that the term proportional to $\partial_r\Delta\phi$ is generally the dominant correction term. This term describes the effect on ν_2 of the fact that the gas falling inward from the clump winds a progressively smaller distance around the star as the clump drifts inward, because the gas falls through a smaller radial distance before colliding with the stellar surface. Hence ν_2 is generally less than ν_{orb} . The fractional decrease in ν_2 is typically $\sim 50\%$ of the fractional increase in ν_1 . The increase in ν_1 and the decrease in ν_2 reduce the frequency separation $\Delta\nu$, which is therefore less than ν_s .

Inward motion of clumps and gas.—The previous analysis shows that $\Delta\nu$ is less than ν_s . A model of the inward drift of clumps near R_{sp} and of the infall of gas from the clumps is required in order to compute the detailed behavior of ν_1 and ν_2 . Here we consider a simple model that captures the basic physics involved.

Let v_{cl} be the inward radial velocity of clumps near R_{sp} , let v_g be the characteristic initial inward radial velocity of the gas stripped from a clump, and assume that $v_{cl} \ll v_g$. Then $\nu_1 \approx \nu_B/(1 - v_{cl}/v_g)$ (see eq. [1]), because the motion near the clump contributes most to Δt . Based on the results of the numerical simulations cited above, we assume that the fractional decrease in ν_2 is equal to one-half the fractional increase in ν_1 . Then $\nu_2 \approx \nu_{orb}(1 - \frac{1}{2}v_{cl}/v_g)$. Our simulations indicate that the initial inward radial velocity of stripped gas is fairly insensitive to the distance of the clump from the star and hence for simplicity we assume that v_g is independent of R_{sp} . We also assume for simplicity that v_{cl} and v_g are approximately constant during the lifetime of a clump.

The inward radial velocity of a clump with angular momentum J caused by removal of its angular momentum by a braking torque N is $v_{cl} = N/(\partial J/\partial r)$. For a clump of mass m in a nearly circular orbit in the Schwarzschild spacetime, $\partial J/\partial r = (m/2)(r - 6M)(r - 3M)^{-3/2}$ and hence

$$v_{cl}(R_{sp}) = \frac{(2/m)(R_{sp} - 3M)^{3/2}}{(R_{sp} - 6M)}N, \quad (3)$$

in units with $G = c = 1$. This expression for v_{cl} diverges at $R_{sp} = 6M$ because the inertia of the clump has been neglected. In reality, $v_{cl}(R_{sp})$ increases as R_{sp} approaches the radius of the ISCO but remains finite (see Fig. 1).

There are several effects that may produce a braking torque N on orbiting clumps. Here we consider one likely possibility. R_{sp} lies inside the star's magnetosphere (MLP98), where the pressure of the stellar magnetic field compresses gas clumps, isolating them from one another. We therefore assume that clumps near R_{sp} are in pressure equilibrium with the stellar magnetic field, that they interact only weakly with each other, and that N is dominated by the interaction of the clump with the stellar magnetic field (see Ghosh & Lamb 1979; Aly & Kuipers 1990; Ghosh & Lamb 1991).




FIG. 1.— Inward radial velocity v_{cl} of clumps orbiting near the sonic radius R_{sp} as a function of R_{sp} , computed numerically (solid lines) and using equation (3) (dotted lines), for (top to bottom) $c_N = 10^{-3}$, 10^{-5} , and 10^{-7} in units of the star's gravitational mass M .

Then $N = \gamma\pi a^2 R_{sp} B_{sp}^2$, where γ is a dimensionless factor where γ is a dimensionless factor ~ 1 that depends on the distortion of the stellar magnetic field by the clump and the clump's shape, a is the mean radius of the clump, and B_{sp} is the magnetic field at R_{sp} . Radiation from the star is likely to maintain clumps at the local Compton temperature, which is almost independent of the distance to the star; we therefore assume that the clump temperature is approximately independent of R_{sp} . For simplicity, we also assume that m is approximately independent of R_{sp} . Then $(3/4\pi)2mk_B T/m_p a^3 = B^2/8\pi = \mu^2 R_{sp}^{-6}$ for a dipolar field with moment μ , where m_p is the mass of a proton, and hence $a \propto R_{sp}^2$ and $N \propto R_{sp}^{-1}$. With these assumptions, the evolution equation for the specific angular momentum u_ϕ of a clump is $du_\phi/d\tau = c_N/r$, where the torque coefficient c_N (which like u_ϕ and the proper time of the clump has the dimensions of a mass) depends on geometrical factors, fundamental constants, and γ . The evolution equation for the radial position of a clump moving in the Schwarzschild spacetime in the absence of significant nongravitational radial forces is

$$\frac{d^2 r}{d\tau^2} = -\frac{M}{r^2} + \left(1 - \frac{3M}{r}\right) \frac{u_\phi^2}{r^3}. \quad (4)$$

We computed $v_{cl}(R_{sp})$ by integrating numerically the clump equations of motion, starting with the values of $\dot{r}$ and $\ddot{r}$ given by equation (3) at large R_{sp} .

Figure 1 shows how v_{cl} varies with R_{sp} for $c_N = 10^{-7}$, 10^{-5} , and 10^{-3} in units of the star's gravitational mass M . These c_N values correspond to clump densities ~ 1 – 10 times the mean density in the disk, for $B_{sp} \sim 10^8$ – 10^9 G. The inward velocity of clumps increases as R_{sp} approaches the ISCO at $6M$, but remains finite. For plausible values of c_N , v_{cl} is small enough even near the ISCO that the inspiral time of the clumps there is long enough to be consistent with the observed coherence of the kilohertz QPOs.

3. RESULTS AND DISCUSSION

The model developed here of the effect of the inward drift of clumps on the frequencies of the kilohertz QPOs has three parameters: the gravitational mass M and spin frequency ν_s




FIG. 2.— Comparison of fits of the SPBF gas dynamical model described in the text (solid curves) with the $\Delta\nu$ - ν_2 correlations observed in four sources. The best-fitting parameter values are Sco X-1: $M = 1.59 M_\odot$, $\nu_s = 352$ Hz, $c_N = 5 \times 10^{-3}$, $\chi^2/\text{dof} = 95.5/46$; 4U 1608-52: $M = 1.76 M_\odot$, $\nu_s = 343$ Hz, $c_N = 2.8 \times 10^{-3}$, $\chi^2/\text{dof} = 16.9/10$; 4U 1728-34: $M = 1.745 M_\odot$, $c_N = 6 \times 10^{-4}$, $\chi^2/\text{dof} = 8.5/6$; and 4U 1820-30: $\nu_s = 279$ Hz, $c_N = 10^{-5}$, $\chi^2/\text{dof} = 7.0/18$. For 4U 1728-34, ν_s was set equal to the 364 Hz frequency of its burst oscillations; for 4U 1820-30, M was set equal to $2.0 M_\odot$, the value indicated by the plateau in its frequency-flux relation (see text).

of the neutron star and the coupling coefficient c_N . If M or ν_s are known, there are only two parameters. Figure 2 shows the best fits of the model to the frequencies of the kilohertz QPOs in Sco X-1, 4U 1608-52, 4U 1728-34, and 4U 1820-30. The model gives good fits (χ^2/dof of 0.4 to 2.1, not including systematic errors), masses (1.59 to $2.0 M_\odot$), spin rates (279 to 364 Hz), and coupling coefficients for all four sources.

The refined version of the SPBF model presented here is consistent with our previous prediction (MLP98) of a plateau in the kilohertz QPO frequency-X-ray luminosity relation when the sonic radius approaches the ISCO. In particular, the fit to the 4U 1820-30 data shown in Figure 2 is consistent with the interpretation that the plateau in the QPO frequency vs. X-ray flux relation reported in this source (Zhang et al. 1998; Kaaret et al. 1999) indicates that (1) there is an ISCO around this neu-

tron star and (2) the asymptotic frequency of the upper kilohertz QPO is the frequency of the ISCO.

Finally, we note that the fundamental frequency in the SPBF model is the orbital frequency at the sonic radius, which is typically $\sim 1,000$ Hz. Hence, even the largest observed difference between $\Delta\nu$ and ν_s (70 Hz in 4U 1728-34) and the largest observed variation of $\Delta\nu$ (100 Hz in 4U 1608-52), although highly significant, represent only very small (5%–7%) deviations of the kilohertz QPO frequencies from the values computed in the circular orbit approximation used previously.

In summary, we have shown that in the sonic-point beat-frequency model, the inward radial motion of the accretion flow causes the frequency separation $\Delta\nu$ between the kilohertz QPOs to be less than the spin frequency ν_s of the neutron star. $\Delta\nu$ is equal to ν_s only if the inflow velocity at the sonic radius is negligible. The inflow velocities expected at R_{sp} naturally produce the small ($\sim 0.2\%$ –5%) corrections to the frequencies predicted for circular orbits (MLP98) needed to explain the differences between $\Delta\nu$ and ν_s and the decreases of $\Delta\nu$ with increasing ν_2 observed in some sources. We have developed a simple physical model of the effect of inflow on the frequencies of the upper and lower kilohertz QPOs. The model fits well the frequencies of the kilohertz QPOs in four Z and atoll sources where they have been fairly accurately measured and gives reasonable values for the masses and spin rates of the neutron stars in these sources.

We emphasize that the frequencies of the kilohertz QPOs predicted by the SPBF model depend sensitively on the properties of the spacetime near the neutron star, including the gravitomagnetic torque produced by the spinning neutron star and whether there is an innermost stable circular orbit. Hence, if the sonic-point model is shown to be correct, measurements of the frequencies of the kilohertz QPOs can be used not only to determine the properties of neutron stars but also to detect and measure gravitational effects in the strong-field regime (see MLP98).

We thank Michiel van der Klis, Mariano Méndez, and Will Zhang for extensive discussions. We gratefully acknowledge the hospitality of the Aspen Institute for Physics, where many of these discussions took place. We are especially grateful to Mariano Méndez for providing the Sco X-1, 4U 1608-34, and 4U 1728-34 data and to Will Zhang for providing the 4U 1820-30 data. This research was supported in part by NSF grant AST 96-18524 and NASA grant NAG 5-8424 at Illinois and NASA grant NAG 5-9756 at Maryland.

REFERENCES

- Alpar, A., & Shaham, J. 1985, *Nature*, 316, 239
 Aly, J.-J., & Kuipers, J. 1990, *A&A*, 227, 473
 Ford, E. C., van der Klis, M., van Paradijs, J., Méndez, M., Wijnands, R. A. D., & Kaaret, P. 1998, *ApJ*, 508, L155
 Ghosh, P., & Lamb, F. K. 1979, *ApJ*, 232, 259
 Ghosh, P., & Lamb, F. K. 1991, in *Neutron Stars: Theory and Observation*, ed. J. Ventura & D. Pines (Dordrecht: Kluwer), 363
 Kaaret, P., Piraino, S., Bloser, P. F., Ford, E. C., Grindlay, J. E., Santangelo, A., Smale, A. P., & Zhang, W. 1999, *ApJ*, 520, L37
 Lamb, F. K., Miller, M. C., & Psaltis, D. 1998, in *The Active X-ray Sky*, ed. L. Scarsi, H. Bradt, P. Giommi, and F. Fiore (Amsterdam: Elsevier), 113 (astro-ph/9802089)
 Lamb, F. K., Shibasaki, N., Alpar, A., & Shaham, J. 1985, *Nature*, 317, 681
 Lorimer, D. R. 2000, to appear in *The Neutron Star – Black Hole Connection*, NATO ASI meeting, Elounda, Crete, 7–18 June 1999 (astro-ph/9911519)
 Méndez, M., & van der Klis, M. 1999, *ApJ*, 517, L71
 Méndez, M., & van der Klis, M. 2000, *MNRAS*, in press (astro-ph/0006243)
 Méndez, M., van der Klis, M., & van Paradijs, J. 1998, *ApJ*, 506, L117
 Méndez, M., van der Klis, M., Wijnands, R. A. D., Ford, E. C., van Paradijs, J., & Vaughan, B. A. 1998, *ApJ*, 505, L23
 Miller, M. C. 1999, *ApJ*, 515, L77
 Miller, M. C. 2000, submitted to *Astr. Lett. Comm.* (astro-ph/0007287)
 Miller, M. C., Lamb, F. K., & Cook, G. B. 1998, *ApJ*, 509, 793
 Miller, M. C., Lamb, F. K., & Psaltis, D. 1998, *ApJ*, 508, 791 (MLP98)
 Psaltis, D. et al. 1998, *ApJ*, 501, L95
 Strohmayer, T. E., & Markwardt, C. B. 1999, *ApJ*, 516, L81
 van der Klis, M. 2000, *ARA&A*, in press (astro-ph/0001167)
 van der Klis, M., Wijnands, R. A. D., Horne, K., & Chen, W. 1997, *ApJ*, 481, L97
 Zhang, W., Smale, A. P., Strohmayer, T. E., & Swank, J. H. 1998, *ApJ*, 500, L171